

For Superior Control of *Rhizoctonia solani*

POTATO SEED-PIECE

MONCOAT[®] MZ seed piece treatment delivers superior long lasting control of both seed-borne and soil-borne diseases to help deliver bigger yields and higher quality potatoes. The active ingredients in Moncoat MZ, flutolanil and mancozeb, provide excellent broad spectrum disease control even under the most severe disease pressure situations.


Key Benefits

- Reliable, effective control of *Rhizoctonia solani*, the pathogen that causes black scurf and stem canker
- Provides added suppression of *Fusarium* dry rot and Silver Scurf
- Helps reduce the spread of seed-borne late blight
- Extended residual disease control
- Dual activity which stops disease progress and prevents future infections
- Systemic activity, providing protection throughout the plant and tuber set
- Alder bark carrier provides improved healing of cut seed pieces
- Excellent yield and quality results

Use Recommendations

- Use Rate 0.75 to 1.0 lb of product per 100 lb of cut seed pieces.
- Apply MONCOAT MZ to seed-pieces immediately after cutting, using dispensing equipment that will give thorough coverage of the dust to the cut seed-piece surfaces.

Rhizoctonia Disease Control 78 Days after Planting



University of Idaho, Dr. Phillip Wharton, Aberdeen, ID 2013 ST = Seed Treatment IF = In-furrow



©2015 Nichino America, Inc. All rights reserved. MONCOAT and Nichino logo are registered trademarks of Nichino America, Inc. Maxim MZ and Quadris are registered trademarks of a Syngenta Group Company. Always read and follow all label directions. 888-740-7700 www.nichino.net MMZ-POT-1501

