

For Control of Broadleaf Weeds

Key Benefits

- Adds power to the tankmix for broad spectrum weed control
- Increases control of hard-to-kill weeds
- Provides strong suppression of small marestail and fleabane
- Reduced risk of volatilization
- Crop safety to young plantings of vines, nut trees, and stone fruit
- 0 day PHI for grape, olive, pome and stone fruit, pomegranate, and tree nuts
- "Caution" label signal word makes Venue easy to use and worker friendly

Use Information

- Can be used in-season, dormant, prebloom, and postharvest on labeled bearing and nonbearing crops.
- Can be used for sucker control on grape, olive, pome and stone fruit, pomegranate, and tree nuts.
- Also labeled for spot treatment and for use in pasture and rangeland, fallow, non-cropland, and noncrop weed control.
- Use 3 to 4 fl oz per acre in a tankmix with glyphosate, 2,4-D, glufosinate, or paraquat, as well as other herbicides.
- Do not use more than 12 fl oz/A per season for all uses.
- Use crop oil concentrate or methylated seed oil to enhance performance.
- Use minimum of 20 gallons of water per acre by ground; 30 to 40 gpa is highly recommended.
- Keep water pH ≤ 7.5
- For best results, spray when weeds are at the rosette stage or 4 inches in height or smaller.
- Venue is a contact herbicide and requires thorough coverage for optimum control.

Venue Controls Tough Broadleaf Weeds Including

- | | | |
|------------------------|------------------------|-------------------|
| ■ Annual sowthistle | ■ Dandelion | ■ Pigweeds |
| ■ Chickweed | ■ Field bindweed | ■ Prickly lettuce |
| ■ Cocklebur | ■ Kochia | ■ Russian Thistle |
| ■ Common lambsquarters | ■ Morningglory species | ■ Stinging Nettle |
| ■ Common mallow | ■ Nightshade species | ■ Yellow Mustard |
| ■ Common purslane | ■ Panicle willowherb | |

Venue Provides Strong Suppression of Small Weeds

- | | |
|----------------|-------------|
| ■ Alkali malva | ■ Fleabane |
| ■ Alkali weed | ■ Marestail |

Control of Silverleaf Nightshade (21 DAT)



Roundup WM 32 fl oz + COC 1%



Venue 4 fl oz +
Roundup WM 32 fl oz + COC 1%

See reverse for additional information >



For Control of Broadleaf Weeds

Mixing Precautions

To avoid incompatibility with glyphosate and paraquat products, follow proper mixing order as directed on Venue product label. Add Venue to the tank first prior to adding any glyphosate or paraquat products. **Do not add Venue to a tank already containing these products.**

Application Precautions for Use in Grapes and Stone Fruit

Venue does not volatilize; however, fine mist droplets can drift and spot foliage and fruit. Avoid spray drifting onto crop. Venue will not translocate into the tree or vine.

Control of Malva and Stinging Nettle (28 DAT)



Untreated



Rely 64 fl oz



Venue 4 fl oz + Rely 64 fl oz

Sucker Control in Grapes (14 DAT)



Untreated Control



Venue 4 fl oz + COC 1%

Control of Panicle Willowweed (3 DAT)



Untreated Control



Venue 3 fl oz + COC 1%



NICHINO
AMERICA®