







PEST MANAGEMENT GUIDE - MELONS FRAC/IRAC Active **Kev Pests** Rates/ REI PHI **Key Benefits Product** Ingredient Controlled Group Acre Unique mode of action for powdery mildew control Powdery 6.0 to 0 **GATTEN®** Flutianil U13 · No cross resistance with existing fungicides 12 hrs mildew 8.0 fl oz Fungicide Day • Translaminar movement – provides protection to both sides of the leaf · Fast knockdown of mites **PORTAL®** · Contact activity on all developmental stages 3 Fenpyroximate 21A of mites - larvae, nymphs, and adults as well Miticide/ Mites 12 hrs 2.0 pints Days Insecticide existing eggs Broad spectrum mite control Active on nymph and adult stages of aphids **Aphids** 2.4 to Superior control of whiteflies (adults and 1st 3.2 fl oz Whiteflies **PQZ®** 1 Pyrifluquinazon 9B 12 hrs instar nymphs) Dav Insecticide · Causes rapid feeding cessation 3.2 fl oz Leafhoppers Contact and translaminar efficacy 14.0 to Halts feeding damage immediately Leafhoppers 21.0 fl oz · Effective on all life stages of insects controlled **TORAC®** Tolfenyprad (eggs, larvae, nymphs, adults) 21A/39 **Aphids** 17.0 to Insecticide · Contact mode-of-action which works by 12 hrs Flea beetle 21.0 fl oz Day inhibiting cellular respiration in the insect Powdery mildew · Added benefit of contact fungicide activity for **Thrips** 21.0 fl oz powdery mildew control and suppression of Downy mildew downy mildew (suppression)

Some products are not registered for use in all states. Consult with Nichino representative or local regulatory agency for registration status in your state.

NICHINO AMERICA