

## CITRUS

BEXAR® CA\* is a broad-spectrum insecticide for control of several key pests in citrus, including citrus thrips. BEXAR CA provides fast and complete control of multiple key pests in citrus with a single application, including those pests known to be resistant to other modes of action.

### Mode of Action

- Mitochondrial Electron Transport Inhibitor (METI) – IRAC Group 21A Insecticide/Miticide
- Disrupts cellular respiration
- Rapid feeding inhibition and mortality within 24 to 48 hours
- Unique mode of action for citrus thrips, soft scales, katydids

### Key Pests Controlled

Citrus thrips, Citricola scale, Barnacle scale, Katydids, Aphids, Asian citrus psyllid (ACP), Bud mite, Citrus mealybug, Brown soft scale

### Key Benefits

- An alternative mode of action for controlling citrus thrips
- Controls insects resistant to other chemistries
- Effective on all life stages of target pests including eggs
- Broad spectrum activity against multiple pests (insects and mites)
- Excellent resistance management tool
- Alternative to organophosphate (OP), carbamate, pyrethroid, neonicotinoid, and spinosyn insecticides
- Not a restricted use product; no foul odor formulation
- Stable formulation compatible with other insecticides, as well as fungicides and fertilizers


### Use Recommendations

Pest	Rate/Acre
Citrus Thrips	27 fl oz/A
Asian citrus psyllid / Katydid	21 to 27 fl oz/A
Citrus bud mite, Citrus mealybug, Soft scale insects (citricola scale, barnacle scale, brown soft scale)	24 to 27 fl oz/A

- Apply by ground with a minimum of 100 gallons of water per acre
- Use an adequate spray volume to achieve complete coverage
- Use 415 oil or equivalent to obtain good coverage

### Product Information

- BEXAR CA is a contact and residual insecticide. Thorough spray coverage is critical to obtain optimal control of the target pest(s)
- Nonselective chemistry; follow use directions carefully regarding application timing and county restrictions
- Reentry Interval (REI): 12 hours
- Preharvest Interval (PHI): 3 days



\*BEXAR CA is for use only in the following counties in California: Fresno, Imperial, Kern, Kings, Madera, Riverside, San Bernardino, San Diego, Tulare, and Ventura counties.

# Efficacy of BEXAR CA on Citrus Thrips

## Citrus Thrips Fruit Scarring




Untreated



BEXAR CA @ 27 fl oz/A

## Commercial Application of BEXAR CA to Control Citrus Thrips in Navel Oranges, Tulare County, CA



Refer to the global MRL database for current established tolerances  
[www.globalmrl.com/db#query](http://www.globalmrl.com/db#query)

